arXiv:0807.2178v1 [cs.CG] 14 Jul 2008

Ranking Unit Squares with Few Visibilities

Bernd Gärtner*

August 4, 2008

Abstract

Given a set of n unit squares in the plane, the goal is to rank them in space in such a way that only few squares see each other vertically. We prove that ranking the squares according to the lexicographic order of their centers results in at most $3n-7$ pairwise visibilities for $n \geq 4$. We also show that this bound is best possible, by exhibiting a set of n squares with at least $3n-7$ pairwise visibilities under any ranking.

1 Problem statement

The unit square with center $c \in \mathbb{R}^2$ is the set

$$S(c) = \{p \in \mathbb{R}^2 \mid \|p-c\|_\infty \leq 1\}.$$

Given a set $\mathcal{S}$ of n unit squares (simply called squares in the sequel), a *ranking* of $\mathcal{S}$ is a sequence $\rho = (S_1, S_2, \ldots, S_n)$ such that $\mathcal{S} = \{S_1, S_2, \ldots, S_n\}$. For $i < k$, S_i *sees* S_k under ρ if there exists a point $p \in \mathbb{R}^2$ such that

$$\begin{array}{lll} p & \in & S_i \cap S_k, \\ p & \notin & S_j, \qquad i < j < k. \end{array}$$

The graph on $\mathcal{S}$ formed by all pairs $\{S_i, S_k\}$ such that S_i sees S_k under ρ is called the *visibility graph* of ρ and will be denoted by $G(\rho)$.

The goal is to find a ranking ρ such that $G(\rho)$ has as few edges as possible. We do not know how to find the best ranking for a given set $\mathcal{S}$, but we prove that there is always a ranking ρ under which $G(\rho)$ has no more than $3n-7$ edges. For some sets $\mathcal{S}$, this is the best bound that can be achieved.

This research is motivated by similar questions for intervals in $\mathbb{R}^1$ [1].

2 Main result

Given a ranking $\rho = (S_1, S_2, \ldots, S_n)$, the center of square S_i will be denoted by $c_i = (x_i, y_i), i = 1, 2, \ldots, n$. We will repeatedly use the following simple fact.

*Institute of Theoretical Computer Science, ETH Zürich, CAB G32.2, CH-8092 Zürich, Switzerland. Email: gaertner@inf.ethz.ch

Observation 1 *Let $\rho = (S_1, S_2, \ldots, S_n)$ be a ranking, and suppose that there are centers c_i, c_j, c_k with $i < j < k$, such that c_j is contained in the (axis-parallel) rectangle spanned by c_i and c_k. Then $\{S_i, S_k\}$ is not an edge of $G(\rho)$.*

Proof. Center c_j being contained in the rectangle spanned by c_i and c_k is easily seen to be equivalent to $S_j \supseteq S_i \cap S_k$. It follows that S_i does not see S_k under ρ. □

The size (number of edges) of the visibility graph may be $\Theta(n^2)$ for a "bad" ranking, see Figure 1 (left). The right part of the figure depicts the lexicographic ranking. According to the next lemma, this ranking always results in $O(n)$ visibilities.



Figure 1: Upward view on a ranked set of squares. Left: All squares in the lower half see all squares in the upper half. Right: the lexicographic ranking incurs only linearly many visibilities.

Lemma 2 *Let $\rho_{lex} = (S_1, S_2, \ldots, S_n)$ be the lexicographic ranking, meaning that $i < j$ if and only if the center of S_i is lexicographically smaller than the center of S_j. Then $G(\rho_{lex})$ is a planar graph. More precisely, the straight-line embedding of $G(\rho_{lex})$ obtained by mapping each square S_i to its center $c_i \in \mathbb{R}^2$ is a plane graph.*

Proof. We show that if two segments $\overline{c_i c_j}$ and $\overline{c_k c_\ell}$ cross, then $G(\rho_{lex})$ contains at most one of the edges $\{S_i, S_j\}$ and $\{S_k, S_\ell\}$.

Since deletion of other squares can only add visibilities between the four squares involved in the crossing, we may w.l.o.g. assume that $\mathcal{S} = \{S_1, S_2, S_3, S_4\}$. By lexicographic order of the centers, there are only two cases.

Case (a): Segments $\overline{c_1 c_3}$ and $\overline{c_2 c_4}$ cross. If y_2 is between y_1 and y_3, we get that c_2 is in the rectangle spanned by c_1 and c_3, so $\{S_1, S_3\}$ is not an edge of $G(\rho_{lex})$ by Observation 1.

If y_2 is not between y_1 and y_3, we either have $y_2 < \min(y_1, y_3)$ and thus $y_4 > y_3$ (otherwise, there would be no crossing), or $y_2 > \max(y_1, y_3)$ and thus

$y_4 < y_3$. In both cases, y_3 is between y_2 and y_4 which means that c_3 is in the rectangle spanned by c_2 and c_4. This in turn shows that $\{S_2, S_4\}$ is not an edge of $G(\rho_{\text{lex}})$.

Case (b): Segments $\overline{c_1c_4}$ and $\overline{c_2c_3}$ cross. An easy case occurs if one of c_2 and c_3 is in the rectangle spanned by c_1 and c_4, since Observation 1 then implies that $\{S_1, S_4\}$ is not an edge.

Otherwise, we have

$$y_2 < \min(y_1, y_4) \leq y_4 \tag{1}$$

and thus

$$y_3 > \max(y_1, y_4) \geq y_1 \tag{2}$$

(because of the crossing), or $y_3 < \min(y_1, y_4)$ and $y_2 > \max(y_1, y_4)$. Let us only treat the first case; the second one is symmetric under exchange of indices 2 and 3.

We now show that every point in $S_1 \cap S_4$ is in $S_2 \cup S_3$, given that $S_2 \cap S_3 \neq \emptyset$. Therefore, if $\{S_2, S_3\}$ is an edge of $G(\rho_{\text{lex}})$, then $\{S_1, S_4\}$ is not an edge.

Let $p = (x, y)$ be any point in $S_1 \cap S_4$. From $x_4 - 1 \leq x \leq x_1 + 1$ and lexicographic order it follows that

$$x_2 - 1, x_3 - 1 \leq x \leq x_2 + 1, x_3 + 1. \tag{3}$$

Using $y_4 - 1 \leq y \leq y_1 + 1$ together with (1) and (2), we also conclude that

$$y_2 - 1 \leq y \leq y_3 + 1. \tag{4}$$

If y in addition satisfies $y \leq y_2 + 1$, (3) and (4) imply that $p \in S_2$. But if $y > y_2 + 1$, we can use the assumption $S_2 \cap S_3 \neq \emptyset$ to conclude that $y_3 - y_2 \leq 2$ and hence $y > y_3 - 1$. With (3) and (4), we then get $p \in S_3$. □

Now we are ready to prove our main theorem.

Theorem 3 *If $n \geq 4$, $G(\rho_{\text{lex}})$ has at most $3n - 7$ edges.*

Proof. An upper bound of $3n - 6$ already follows from Lemma 2. In order for this bound to be tight, the outer face of $G(\rho_{\text{lex}})$'s straight-line embedding would have to be a triangle Δ spanned by three centers c_i, c_j, c_k, $i < j < k$, and with all other centers inside Δ. Indeed, for $n = 3$ this is possible, but for $n \geq 4$, we get a contradiction: Let $c = (x, y)$ be any center distinct from c_i, c_j, c_k. From $c \in \Delta$, it easily follows that $c_i < c < c_k$ (comparison being lexicographically).

If y is between y_i and y_k, c is in the rectangle spanned by c_i and c_k, so $\{S_i, S_k\}$ can't be an edge of $G(\rho_{\text{lex}})$ by Observation 1.

If y is not between y_i and y_k, then y must be between y_i and y_j, *and* between y_j and y_k. Depending on whether $c_i < c < c_j$ or $c_j < c < c_k$, we get that either $\{S_i, S_j\}$ or $\{S_j, S_k\}$ is not an edge of $G(\rho_{\text{lex}})$. □

3 Lower Bound

The bound of $3n - 7$ derived in Theorem 3 is best possible in the worst case, not only under the lexicographic ranking, but under *every* ranking. For a proof by picture see Figure 2.



Figure 2: A set of n unit squares such that the visibility graph $G(\rho)$ has at least $3n - 7$ edges for every ranking ρ: a square from the middle bunch of $n-2$ squares always sees the next square above it in the bunch (these are $n-3$ visibilities), and it sees (or is seen by) both of the two special squares ($2n - 4$ visibilities).

Acknowledgment

The problem of ranking unit squares was given to me as an undergraduate student by Emo Welzl in early 1989. After some initial programming tasks, this was the first actual research assignment during my time as "Forschungstutor" (research assistant) with Emo.

My initial manuscript from 1989, typeset in bumpy LaTeX and containing manually drawn figures, is lost. But throughout the almost twenty years that have passed since then, I could never forget this first (simple) result of mine.

While writing it up now, many memories of all these years have come back to me with surprising strength. I thank Emo for bringing the problem to my attention, and for so much more.

References

[1] Herbert Edelsbrunner, Mark H. Overmars, Emo Welzl, Irith Ben-Arroyo Hartman, and Jack A. Feldmann. Ranking intervals under visibility constraints. *International Journal of Computer Mathematics*, 34:129–144, 1990.